

**Ndukwu CI
Egbuonu I
Ulasi TO**

Nutritional status and sociodemographic characteristics of ‘urban poor’ school children in Onitsha, Southeast Nigeria

DOI:<http://dx.doi.org/10.4314/njp.v41i3.5>

Accepted: 11th February 2014

Ndukwu CI (✉)
Egbuonu I, Ulasi TO
Department of Paediatrics
Nnamdi Azikiwe University Teaching
Hospital Nnewi.
Anambra State, Nigeria.
Email: ifeyc@yahoo.com

Abstract Background: The neglect of the health and nutritional status of school aged children has adverse consequences on their long term cognition and survival. Sequel to emerging issues on urbanization and the health of school children, this study focused on the nutritional status and sociodemographic characteristics of ‘urban poor’ school children in Onitsha, a major city in South East Nigeria.

Objectives: The study was aimed at determining the nutritional status and evaluating the effect of certain sociodemographic factors on the nutritional status of ‘urban poor’ public primary school children in Onitsha, South East Nigeria

Method: This was a cross sectional study of 788 children aged 6 to 12 years, randomly selected from 12 public primary schools in Onitsha metropolis. Their anthropometric measures were used to determine their nutritional status. Data on their sociodemographic characteristics was obtained from their parents or caregivers using

interviewer administered standardized questionnaires. Analysis was done using Microsoft Office Excel 2007, SPSS version 17 and Epi-info version 3.5.1 software packages

Results: Stunting was the predominant form of under nutrition with a higher prevalence in the slum resident children. Greater than 60% of all the children studied were from large families with more than 4 children. Family size, however, age and gender were not significantly associated with the nutritional status of the children. Relatively though, a greater percentage of the stunted children were from large families.

Conclusion: The nutritional status of ‘urban poor’ school children especially in slums in Onitsha is suboptimal compared with their counterparts in developed countries of the world. There is need for regional studies of children with similar characteristics who should be considered when favourable health policies are being made for children.

Introduction

The nutritional status of children plays a central role in their health, with malnutrition accounting for 45% of all deaths among children under the age of five and stunting growth among a further 165 million¹. With the progressive success of various child survival programmes, more children are reaching school age (ages 6 to 12)². Unfortunately, minimum attention has been paid to the health and nutritional status of the school aged child, especially in developing countries with more emphasis laid on improving their educational access³. However, undernourished and unhealthy school children will fail to reach their full cognitive potential with ultimate reduction in intellectual achievement in school, and poor work capacity later on in adulthood^{3,4,5}.

Hundreds of millions of children in the world’s urban areas are growing up amidst scarcity and deprivation with increasing incidence of childhood malnutrition and its complications^{4,6}. Certain sociodemographic characteristics like age, gender and birth order have also been found to influence their health and nutritional status^{7,8,9}. There is need to study the prevailing characteristics in different areas so as to identify those that need specific interventions.

This cross sectional study is focused on ‘urban poor’ public primary school children residing in slum and non-slum areas of Onitsha, Anambra state with the aim of drawing attention to their needs. Not all of the urban poor live in slums, and not every inhabitant of a slum is poor⁴. However, most of the poor are known to attend

the public schools that are 'tuition free' in the state while the 'well off' send their children to fee paying private institutions¹⁰. The results from this study will provide evidence on school-age malnutrition in this part of the country, drawing attention to the need for effective interventional strategies that will improve the health outcome of these children.

Method

This was a cross sectional study conducted in Onitsha from September to November, 2010. Onitsha is a riverine port on the Eastern bank of the River Niger. It is the most rapidly expanding commercial centre in Eastern Nigeria¹¹. Densely populated with four recognized slum areas, major parts lack basic social services¹¹.

Children of school age in public schools in the slums and non slum areas formed the study population. Twelve schools were randomly selected for the purpose of this study; six were located in the slum areas making up approximately 30% of all the public primary schools in the slum areas while the corresponding numbers of public schools were located in the non slum areas. The subjects were selected by the stratified multistage random method. A sample size of 384 was determined using a standard formula¹².

Approval for this study was obtained from the Ethical Committee of Nnamdi Azikiwe University Teaching Hospital as well as the Anambra state Universal Basic Education Board. Children were recruited prorata based on the school population and their age and sex distribution. Consenting parents or guardians were invited to the schools to provide answers to interviewer-administered questionnaires on the demographic characteristics of the participants. Excluded from the study were those whose parents declined consent and those who had obvious chronic ailments. The children's weights and heights were subsequently measured using standard methods. A 'Hana' bathroom scale, model BR 9011, ISO 9001:2000 certified by SGS, was used to measure the weights of the children, dressed in lightweight gowns or pants. The weight was recorded to the nearest 0.5kilogramme. The scale needle was returned to zero before daily weighing and thereafter, a known weight was placed to assess the reliability of the scale. The heights were measured to the nearest 0.5 centimetre with a special locally constructed wooden stadiometer¹³. Random re-measuring of the weights and heights of some of the children was done to assure quality.

Data analysis was done using Microsoft Office Excel 2007, SPSS version 17 and Epi-info version 3.5.1 software packages. The statistical tests were carried out at significance (p value) of less than 0.05. The z-scores for the anthropometric indices were obtained using WHO/Z-score reference (2007), and the WHO/NCHS 1978 international reference standards (Epi-nut from epi-info version 3.5.1). The anthropometric indices were com-

pared with the median for the age according to the 2007 WHO charts.

Result

Complete data and measurements were obtained from 788 children.

Demographic characteristics

The children studied included 398 males and 390 females. Male: female ratio was 1.02:1 and the mean age of the respondents was 8.9±1.9years. (Table 1) Ten year olds accounted for the greatest percentage (22.3%) of children recruited. Table 2 shows the sociodemographic characteristics of the children studied in the slums and non-slum areas. Most of the respondents were from monogamous families and from families with more than four children. (69% in the slums and 64% in the non-slums).

Table 1: Age and Sex distribution of the study population per area

Age (in years)	Slum Male n=398 (%)	Non slum Female n=390(%)	Total n=788(%)
6	62 (15.6)	62 (15.9)	124(15.7)
7	44 (11.1)	46 (11.8)	90(11.4)
8	62 (15.6)	64 (16.4)	126(16.0)
9	48 (12.1)	44 (11.3)	92(11.7)
10	88 (22.1)	88 (22.6)	176(22.3)
11	60 (15.1)	50 (12.8)	110(14.0)
12	34 (8.5)	36 (9.2)	70(8.9)
Mean Age ±SD	8.9±1.9	8.9±1.9	

Table 2: Sociodemographic characteristics of the children

Demographic characteristics	Slum group n=394 (%)	Non-slum group n=394 (%)	X ²	p-value
<i>Family setting</i>				
Monogamy	365 (92.6)	357 (90.6)	1.29	0.525
Polygamy	22 (5.6)	26 (6.6)		
Others*	7 (1.8)	11 (2.8)		
<i>Number of children</i>				
≤ 4	118 (30)	130 (33)	2.42	0.120
> 4	272(69)	253(64.2)	df=1	
DNK	4(1.0)	11(2.8)		
<i>Birth Order</i>				
1 st	66(16.8)	81(20.6)	2.41	0.299
2 nd	64(16.2)	67(17.0)		
3 rd	75(19.0)	69(17.5)		
4 th	63(16.0)	52(13.2)		
>4	118(30.0)	110 (27.9)		
DNK	8 (2.0)	15 (3.8)		
<i>Ethnic group</i>				
Ibos	360 (91.4)	388 (98.5)	20.65	0.000
Non Ibos	34 (8.6)	6 (1.5)		

*Others - widowed, divorced, separated

Nutritional Status and association with sociodemographic factors

Stunting was the predominant form of under nutrition in the children with significantly higher prevalence in the slum than in the non slum areas. Table 3 shows the prevalence and patterns of under nutrition in all the children. Prevalence of underweight was 11.5% in the slum and 6.3% in the non slum area ($X^2 = 4.85$, $p=0.028$), and for stunting was 16.8% in the slum compared to 6.6% in the non slum area ($X^2 = 19.81$, $p=0.000$).

Nutritional status	Slum n (%)	Non slum	X^2	P-value
Underweight (6-10years) n=304(%)	35 (11.5)	20 (6.3)	4.85	0.028
Stunting (6-12 years) n=394 (%)	66 (16.8)	27(6.9)	19.81	0.000
Wasting (6 -10 years) n=304(%)	4 (1.3)	2 (0.7)	0.67	F=0.685
	299 (98.7)	301(99.3)		

Six year old males had the highest prevalence of underweight. The mean age of the underweight children was less than the mean age of the normal. Ten year olds in the slums (28.8%) and 12 year olds in the non-slums (29.6%) had the highest prevalence of stunting. More males were stunted than females though not significantly. Only one female child was overweight in this study.

Family size was not significantly associated with the nutritional status of the children Table 4. Relatively, of all from families with less than 4 children, only 11.9% were stunted as against 18% of children from the families with more than 4 children in the slums (not represented). In the non-slum areas, 6.2% of all from families with less than 4 children were stunted compared with 6.7% amongst those from families with more than 4 children. No significant association was equally demonstrated between the birth order of the children and their nutritional status.

Table 4: Family size, birth order and Stunting in the Children
Mean anthropometric measures in both localities

Demo-graphic factor	Stunting		X^2	P-value	Stunting		X^2	P-value
	Slum	Non-Slum			Slum	Non-Slum		
	Yes n=66 (%)	No n=328 (%)			Yes n=27 (%)	No n=367 (%)		
<i>Family size</i>								
≤4	14 (21.2)	104 (31.7)	3.46 df=2	0.1 77	8(29.6)	122 (33.2)	1.08 df=1	0.298
>4	49 (74.2)	223 (68)			17(63)	236 (64.3)		
DNK	3 (4.6)	1(0.3)			2 (7.4)	9(2.5)		
<i>Birth order</i>								
1 st to 4 th	41 (62.1)	226 (68.9)	1.16 df=1	0.2 82	14 (51.9)	254 (69.2)	3.48 df=1	0.61
> 4 th	22 (33.4)	97 (29.6)			11 (40.7)	100 (27.2)		
DNK	3 (4.5)	5 (1.5)			2 (7.4)	13 (3.5)		

There was a significant difference between the mean weight and height of children in both communities. (Table 5) Children from the non-slum area were significantly heavier and taller than their slum counterparts. Male school children from the slums were significantly shorter than their counterparts in the non-slum areas ($t=3.09$, $p=0.002$). The mean weights and heights for children of same age and gender from the 2007 WHO charts were all greater than the means from the study population. Figures 1 and 2 are illustrations of approximate mean weights and heights of the children according to gender and area of residence, and compared with WHO mean weights and heights for children of same age and gender.

Table 5: Mean anthropometric parameters by age and gender in both groups

Characteristic	Study group n=394 Mean ± SD	Control group n=394 Mean ±SD	t-test	P-value	
Mean Weight (kg)	26.9±6.2	28.1±7.3	2.39	0.017	
Mean Height(cm)	129.6±10.7	132.4±11.2	3.50	0.000	
Age	Sex				
6 years	Female	Weight 19.6±2.5 Height 116.1± 6.5	20.3±3.3 117.4± 5.7	0.958 0.811	0.342 0.421
	Male	Weight 19.7±2.8 Height 113.9±4.5	20.5±2.1 117.4±5.3	1.22 2.79	0.228 0.007
7 years	Female	Weight 22.2±2.5 Height 122.3±5.7	22.4±2.2 122.0±4.7	0.284 0.337	0.777 0.822
	Male	Weight 21.6± 2.7 Height 120.1 ±6.2	22.7± 2.6 122.4±4.4	1.42 1.49	0.163 0.143
	Female	Weight 24.8±3.1 Height 127.8±5.5	24.9±3.2 127.8±5.9	0.218 0.55	0.827 0.957
8 years	Male	Weight 24.5±2.6 Height 126.1±5.5	25.4 ±2.4 129.0±4.4	1.40 2.29	0.165 0.026
	Female	Weight 26.9±3.6 Height 131.1±5.0	27.4±3.4 132.1±5.8	0.473 0.575	0.638 0.568
	Male	Weight 26.6±2.4 Height 129.8±6.2	28.4±4.5 134.3±6.2	1.70 2.48	0.950 0.017
9 years	Female	Weight 28.3±3.2 Height 133.7±5.3	32.2±8.4 139.5±6.8	2.90 4.51	0.005 0.000
	Male	Weight 29.3±4.5 Height 134.2±6.2	30.6±4.6 137.3±7.3	1.31 2.15	0.194 0.035
	Female	Weight 33.4±4.7 Height 140.7±8.0	32.6±4.6 140.4±7.0	0.564 0.142	0.575 0.889
10 years	Male	Weight 31.6±3.8 Height 136.8±6.6	31.8±5.0 140.3±7.0	0.189 2.01	0.851 0.049
	Female	Weight 37.5±5.3 Height 143.5±7.8	40.1±8.7 149.2±6.3	1.09 2.39	0.281 0.022
	Male	Weight 35.6±4.1 Height 143.2±4.8	36.5±5.5 144.1±6.3	0.496 0.459	0.623 0.649

Fig 1: Mean weights of girls (per age group) in study areas and mean from WHO (2007 charts)





Fig 2: Mean heights of boys (per age group) in study areas and mean from WHO (2007 charts)



Discussion

The nutritional status of school age children in public primary schools in Onitsha is sub-optimal and concurs with the pattern in national studies of the nutritional status of Nigerian children.^{13,14,15} Similar to findings from these studies, stunting had the highest prevalence, followed by underweight suggesting that these children have been exposed to long standing conditions that are inimical to good health and nutrition.

The prevalence of underweight and stunting in the slums of 11.5% and 16.8%, respectively are similar though marginally higher than the results of the 2004 Nigerian Demographic and health survey Ed Data Survey (the most recent National survey focused on school-age children) of 11% underweight and 13% stunting amongst school age children from South East Nigeria, which is the study location.¹⁴ The Nigerian economy has been retrogressive in the past decade with increasing number of hungry, extremely poor people, which may be partly responsible for the non improvement in the nutritional status in this index study compared with the National study conducted 6years earlier. The lower prevalence of 6.3% underweight and 6.9% stunting in the non-slum areas suggests interplay of factors like the environment and abject poverty in the slum areas. The higher prevalence of under nutrition in high density areas in other parts of the country may be due to other factors that affect the nutritional status of children such as the educational status of the parents as well as other socioeconomic and environmental factors that need to be identified.^{13,15,16,17} The two groups studied were selected from public, non-fee paying primary schools with the aim of assessing children from the lower social strata of the society. However, an assessment of the social class showed that more children from the slum areas were from the lower strata. This may be responsible for the greater prevalence of stunting in that group.

The prevalence of under nutrition in this study sharply contrasts with the 2.7% reported for school-aged children in the United States of America.¹⁸ The disparity is not surprising as under nutrition is rooted in poverty.

The index study was carried out in a Sub Saharan developing country characterized by widespread poverty with inequitable distribution of wealth and non-existence of a social security system. The United States, a Western developed country, has existing policies that result in social and health stability. Concomitantly, the mean anthropometric indices in the urban poor children were all low compared with the value from the WHO growth charts for corresponding age and gender. The charts were modified from the 1977 NCHS/WHO growth curves derived from a healthy, non-obese Caucasian population.¹⁹ It is disheartening that these children are still disadvantaged more than three decades later.

Over nutrition is not a problem amongst the urban poor in this part of Nigeria, contrary to findings in the developed world where the prevalence has shown a steady rise.^{20,21} This is expected as the Western diets of high saturated fats, sugars, and refined food is expensive and not readily available to the poor in this part of the world unlike in the developed countries. Studies on Nigerian children have equally reported an increased prevalence of over nutrition in children from the higher socioeconomic classes.^{22,23}

Some sociodemographic characteristics contributed to the nutritional status of the study population. The mean age of those who were underweight was lower than that of the general population possibly because older children may be self sufficient, resorting to other measures such as begging, performing menial jobs and scavenging to satisfy their hunger. The higher mean age for the stunted children corresponds with other findings that have demonstrated the 'progression of height deficit' with the increasing age of children. Stunting is a reflection of longstanding under nutrition occurring from a cumulative process of pre-natal and early childhood malnutrition and stunted children are likely to remain stunted even into adulthood if they remain in a deprived environment.²⁴

That males were generally most affected by under nutrition is similar to other findings in Nigeria and other parts of the world.^{13,14,24} This may be attributable to the later onset of pubertal growth spurt in males and the findings that males usually start life at a disadvantage sustained even up to school age.²⁵ Uthman found a higher prevalence of stunting and underweight among males with gender having the strongest independent effect on the risk of stunting.¹⁷ A contradicting finding in India was postulated to be due to gender bias and parental preferences for male children in their society.⁹

Poorer nutritional status has been documented in children of higher birth order and from larger household size.^{8,9} The relative association noted with twice the percentage from bigger family size being stunted in this study is expected because with more mouths to feed, the individuals may likely get smaller portions. In India equally, children from joint families were found to be more likely to suffer chronic malnutrition than those from nuclear families which are smaller in size.⁹

Conclusion

The nutritional status of 'urban poor' school children in this study is suboptimal and associated with certain demographic characteristics like family size, age and gender. It is evident however, that there may be some interplay of other factors especially the social class and certain environmental characteristics, which may be why the slum resident children had worse nutritional status. There is need for further studies focusing on the social classification in terms of the educational level of the parents, their occupation and income, and environmental characteristics to get a holistic picture of the reasons for the prevalent malnutrition. It is also recommended that a school-based feeding programme should be introduced in public primary schools in the area of study.

Conflict of Interest : None

Funding: None

Acknowledgements

We are grateful to Dr Onyeonoro, a public health physician, Mr Kenechukwu Odumodu, an accountant with Schlumberger company, Ghana and Engr. Ikechukwu Odumodu, an author and director of a private school in Nigeria, who helped with different areas of the analysis. Dr Ndukwu Chibuzo is greatly acknowledged for his support during the entire study.

References

1. The daily impact. Global health news and commentary: Lancet series shows malnutrition causes 3.1 million child deaths. June 6, 2013.
2. Drake L, Maier C, Jukes M, Patrikios A. School-Age Children: Their Nutrition and Health.
3. In: Partnership for child development. School-Age Children, Their Nutrition and Health. 2002; 5
4. Gunde R. School Children in the Developing World: Health, Nutrition and School Performance. In a 2-day International Workshop under the Global Impact Research Initiative. The Regents of the University of California 2009.
5. UNICEF. The State of the World's Children. New York: Oxford University Press; 2012.
6. De Onis M, Frongillo EA, Blossner M. Is malnutrition declining? An analysis of changes in levels of child malnutrition since 1980. *Bull WHO* 2000; 78: 1222-1223.
7. Environmental Health Project. Improving the health of the urban poor Learning from USAID experience. *Strategic Report 12*. 2004; 1-67.
8. Akor F, Okolo S, Okolo A. Nutritional status of newly enrolled primary school children in Jos-Plateau, Nigeria. *Pak J Nutr* 2010. 9(12) :1166-70.
9. Mushtaq MU, Gull S, Khurshid U, Shahid U. Prevalence and socio-demographic correlates of stunting and thinness among Pakistani primary school children. *BMC public health* 2011, 11: 790 doi:10.1186/1471-2458-11-790.
10. Srivastava A, Mahmood SE, Srivastata PM, Shrotriya VP, Kumar B. Nutritional status of school-age children - A scenario of urban slums in India. *Archives of public health* 2012;70:8 doi 10.1186/0778-7367-70-8
11. Olaniyan DA, Obadara OE. A critical review of management of primary education in Nigeria. *Int J Afr Am Studies* 2008; 8(1):19
12. Ajayi MO, Ejindu CF, Okojie CC, Udeaja F, Onweluzo F. Structure Plan for Onitsha and Satellite towns. United Nations Human Settlements Programme (UN-HABITAT), 2009 ;14-30.
13. Araoye MO. Subjects Selection. In: Araoye MO(Ed). *Research Methodology with statistics for Health and Social Sciences*. Ilorin: Nathadex (Publishers). 2003; 116-29.
14. Ajayi IO, Akinyinka OO. Evaluation of the nutritional status of first year school children in Ibadan, Southwest Nigeria. *Afr. J. Med.Sci.* 1999; 28: 59-63.
15. National Population Commission (NPC) [Nigeria] and ORC Macro .Nigerian Demographic and health survey EdData Survey 2004. Calverton, Maryland: National Populaton Commission and ORC Macro.2005; 3-40.
16. Amuta EU, Houmsou RS. Assessment of Nutritional Status of School Children in Makurdi, Benue State. *Pak J Nutr* 2009; 8: 691-694.
17. Abidoya RO, Ihebuzor NN. Assessment of nutritional status using anthropometric methods on 1-4 year old children in an urban ghetto in Lagos, Nig. *Nutr Health* 2001; 15: 29-39.
18. Uthman O.A. Using extended concentration achievement indices to study socioeconomic inequality in childhood malnutrition: the case of Nigeria. *Int J Equity* 2009; 8 .
19. Fryar CD, Ogden CL. Prevalence of underweight among children and adolescents: United States, 2003-2006. *National Health and Nutrition Examination Survey*. Health E-stats. 2009.
20. De Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a
21. WHO growth reference for school-aged children and adolescents. *Bull WHO* 2007; 9: 649-732.
22. Cynthia O, Margaret C. Prevalence of obesity among children and adolescents: United States, Trends 1963-1965 Through 2007-2008. Centers for Disease Control and Prevention. National Center for Health Statistics E-stat.
23. Sharma AJ, Grummer-Strawn LM, Dalenius K, Galuska D, Anandappa M et al. Obesity Prevalence among low-income preschool-aged children- United States, 1998-2008. Centers for Disease Control and Prevention. *Morbidity and Mortality Weekly Report* 2009; 769-73.
24. Owa JA, Adejuyigbe O. Fat mass, fat mass percentage, body mass index and mid-upper arm circumference in a healthy population of Nigerian children. *J Trop Paediat* 1997; 43: 13-19.

-
25. Senbanjo IO, Adejuyigbe O. Prevalence of overweight and obesity in Nigerian pre-school children. *Nutr Health*, 2007; 18: 391-99.
 26. Al-Saffar AJ. Stunting among primary-school children: a sample from Baghdad, Iraq. *East Mediterr Health J*. 2009; 15: No.2.
 27. Akinyinka OO, Sanni KA, Falade AG, Akindele MO, Sowumi A. Arm area measurements as indices of nutritional reserves and body water in African newborns. *Afr. J. Med Med. Sci.* 1999; 28: 5-8.