

Ogunlesi TA

CC –BY A survey of Paediatricians on evidence-based medical practice in Nigeria

DOI:<http://dx.doi.org/10.4314/njp.v45i3.5>

Accepted: 17th September 2018

Ogunlesi TA (✉)
Department of Paediatrics,
Olabisi Onabanjo University
Teaching Hospital,
P. O. Box 652, Sagamu-121001NG,
Ogun State, Nigeria
Email:
tinuade_ogunlesi@yahoo.co.uk

Abstract: *Background:* The increasing need for a uniform standard of medical practice necessitates an examination of the impediments to the use of evidence-based medicine.

Objective: To examine the knowledge, perceived barriers to the use of evidence-based medicine and the associated profession-related factors among paediatricians in Nigeria.

Methods: A cross-sectional survey of paediatricians in Nigeria was conducted between March and April 2013 by Email. The knowledge about evidence-based medicine and systematic reviews was tested using a structured questionnaire and the responses were provided on three-item Likert scale.

Results: The response rate in the survey was 56.5% (65/115). Almost 37% of the participants had postgraduate qualification in or before year 1999 (Group I). Majority (93.8%) of the respondents believed that EBM should form the basis for decision making in

clinical practice while 53.9% believed that EBM is more suitable for the developed world. The overall mean knowledge score was 77.6%. Knowledge about EBM was rated high among 80% of the participants and the proportion with high knowledge was not significantly associated with EBM use in routine practice or prior EBM training. Some of the perceived barriers to EBM use included lack of skills for understanding statistics (81.6%), lack of incentives for the use of EBM (64.7%), lack of time (44.6%) and lack of conclusive evidences (44.6%).

Conclusion: The knowledge of EBM among Nigerian paediatricians was good but a high proportion regarded lack of skills to perform statistical analysis required for EBM as a major barrier to the practice of EBM.

Key words: Evidence-based medicine, Systematic review, Research, Paediatricians.

Introduction

The prime role of research in driving clinical practice has not changed over the years but the application of research in guiding clinical practice is globally getting focused on the critical need for evidences guiding every aspect of decision taking in medical practice.¹ The contents of textbooks, clinical observations and opinions passed down from senior colleagues have hitherto, formed the bedrock of guidance for making clinical decisions.² With the increasing use of evidence-based medicine, clinical decisions are gradually shifting towards the provisions of management protocols and treatment guidelines rather than textbook recommendations and opinions formed in the course of training and practice. In brief, evidence-based medicine strictly entails adherence to the current best practices for the utmost benefit of the patient. This is necessitated by the need for a more guided approach rather than the haphazard

access to a large volume of pockets of research outputs from various parts of the world, some with conflicting reports and some with inadequate information to guide practice.^{2,3} Therefore, evidence-based medicine appears to be the most efficient way of delivering clinical care in the context of rapidly changing trends in medical practice.

At the tertiary level of care, the tripod of clinical duties includes service, teaching and research. A previous study conducted among medical specialists in Nigeria⁴ showed that research function was ranked third by 64.7% of the participants in that study, behind service and training. A similar study revealed that the spectrum of research tilted more towards low-budget cross-sectional surveys and retrospective studies due to the challenges of lack of funds, work overload, time constraints and insufficient infrastructural supports.⁵ In the same study, 37.5% of specialists had never been in-

volved in clinical trials. Incidentally, randomized clinical trials, which represent the peak of clinical evidence through the benefits of elimination of bias, form the core of evidence-based medicine.¹

Evidence-based medicine entails the identification of an intervention or relationship of interest, gathering relevant high-quality randomized controlled trials and conducting statistical analysis, using meta-analysis, to assess the quality of evidence in support of specific outcome variables of interest.^{1,6} This is the basis of the use of various treatment guidelines and management protocols for clinical decision in various parts of the developed world. Commonly accessed resources for evidence-based medicine include the Cochrane Library and the World Health Organisation Reproductive Health Library. These are the commonly used databases which provide information on virtually all aspects of medicine. Of utmost relevance to the practice of paediatrics are the Neonatal and the Infectious Diseases Review Groups which presently contains hundreds of systematic reviews.

With the drive to develop treatment protocols for common paediatric disorders in Nigeria, it is essential to examine the acceptability and understanding of the concept of evidence-based medicine among paediatricians. Therefore, the objective of this study was to examine the knowledge, perceived barriers to the use of evidence-based medicine and the associated profession-related factors among paediatricians in Nigeria.

Materials and Methods

This was a cross-sectional questionnaire-based survey of paediatricians who practiced in Nigeria. The study was carried out electronically using email-distribution of questionnaire to paediatricians on a mailing list. The research was conducted in agreement with the Helsinki Declaration for Human subjects research. The inclusion criteria included Postgraduate Fellowship in Paediatrics obtained from either the West African College of Physicians or the National Postgraduate Medical College of Nigeria and employment as a Consultant Paediatrician within Nigeria. The study was carried out between March and April 2013.

The minimum sample size was determined using the formula; $n = (z^2 \times p \times q) / d^2$ where $z = 1.96$, $p = 0.05$ (proportion of respondents in a similar study who used EBM to take clinical decisions⁷), $q = 0.95$ and $d = 0.05$. The calculated minimum sample size was 72 but additional 50% (36) was added to increase the strength of the study. Therefore, the final sample size was 104.

Out of the 426 names on the mailing list of paediatricians, 78 duplicated names, names with incomplete data (such as place of practice and year of postgraduate qualification) and names of deceased paediatricians were removed, leaving 348 names. Thereafter, the 348 names were arranged alphabetically and systematic random sampling was done using a sampling interval of three

resulting in 115 rather than the targeted 104. The selected 115 paediatricians were enrolled into survey and were contacted via email. The email contained a statement of introduction of the study as well as the request for consent. The email message specified that response to the survey meant consent for enrolment into the study and anyone not willing to participate in the study was allowed to disregard the request. The data collection was carried out between 3rd March and 5th April 2013.

The research tool was a self-designed close ended questionnaire with three sections using information gathered from the Cochrane Handbook for Systematic Reviews of Interventions as a template in most cases.⁸ The first section obtained data on professional parameters such as the year of postgraduate qualification, location of practice in terms of geopolitical zone, sector of practice (public or private), setting of practice (academic or non-academic), current use of EBM in clinical practice and history of previous formal training in EBM. The second section tested the knowledge of participants about EBM using general statements covering a wide scope of the principles and tenets of EBM as well as test of understanding and ability to teach specific statistical items. The third section assessed the participants' attitude to perceived barriers to the use of EBM in clinical practice. Some of the statements were framed in the positive context and others were framed in the negative context and the responses were obtained using a three-item Likert Scale – Agreed, Undecided and Disagreed. The responses to each statement were uniformly assessed without weighting; they were scored 3, 2 and 1 depending on the context of positivity or negativity; the option of "agreed" for a negatively-framed statement earned a score of "1" just as a response of "disagreed" for a negatively-framed statement earned a score of "3" and *vice versa*. For each participant, the total score was converted to percentage based on the number of statements responded to. Knowledge scores of 75% and above were classified as "high" while scores less than 75% were classified as "low" scores. The mean percentage knowledge score was determined for each group of participants.

Data management

Only completed questionnaires returned by email were pooled for analysis using a spreadsheet created with the Microsoft Excel software. Descriptive and inferential statistics were conducted using the SPSS version 20.0 software. Hypotheses were tested using the Chi Square test with either Yate's correction or the Fisher's Exact test as necessary, for proportions of categorical variables and the Student's t-test for the means (\pm Standard deviations) of continuous variables. Professional characteristics of the respondents (year of postgraduate qualification, sector of practice, setting of practice, use of EBM in clinical practice and training in EBM) were related to the knowledge score as well as the perceived barriers to EBM. *P* values less than 0.05 were accepted as statistical significance.

Results

General description of the respondents

Sixty-five out of the 115 participants returned completely filled questionnaire giving a response rate of 56.5%. The 65 paediatricians were distributed across the geo-political zones of the country as follows: 24 (36.9%) from the south-west, 16 (24.6%) from the south-east, 10 (15.5%) from the south-south, 6 (9.2%) each from the north-west and north-central and 3 (4.6%) from the north-east.

The distribution of the respondents according to the year of post-graduate qualification was as follows: 4 (6.1%), 20 (30.8%), 30 (46.2%) and 11 (16.9%) for 1980-1989, 1990-1999, 2000-2009 and ≥ 2010 respectively. They were re-grouped into two as 1980-1999 (Group I) and 2000 and above (Group II) translating to 24 (36.9%) in Group I and 41 (63.1%) in Group II. The major subspecialties included neonatology (17; 26.2%), cardiology (9,13.9%), haematology and nephrology (7; 10.8% each), endocrinology, infectious diseases and ambulatory paediatrics (6; 9.2% each). The remaining 7 (10.7%) included neurodevelopmental paediatrics, gastroenterology and nutrition and respiratory.

Most of the respondents (53; 81.5%) practiced in the public sector; 33 (50.8%) and 29 (44.6%) practiced in Teaching Hospital and specialist hospitals (including Federal Medical Centres) respectively. The settings of practice were sub-classified into two: academic (33; 50.8%) and non-academic (32; 49.2%). Fifty-two (80.0%) were involved in undergraduate training while all the respondents had experience with postgraduate training. Fifty-nine (90.8%) respondents used evidence-based medicine in their routine clinical practice but only 24 (36.9%) actually had training in evidence-based medicine. While all the respondents in Group I ($n = 24$) used EBM in clinical practice, 35 (85.4%) of Group II used EBM in clinical practice (Fisher's Exact Test = 5.872; $p = 0.07$). A significantly larger proportion of respondents in Group I (17; 70.8%) received training on EBM compared to respondents in Group II (7; 17.1%) ($\chi^2 = 18.786$; $p < 0.001$).

General knowledge and attitude of respondents about EBM

As shown in Table 1, all the respondents agreed that current research findings are useful in routine care of patients, 76.9% craved for further use of EBM in their daily practices while 92.3% were willing to learn the skills required to incorporate EBM into their routine practices. However, 86.1% believed EBM is patient-centred, improves the quality of care available to patients (89.3%), should form the basis of decision making in clinical practice (93.8%) and should improve the design of medical curriculum (83.1%). On the other hand, the respondents identified drawbacks such as EBM not taking into consideration the limitations in clinical practice (55.4%) and EBM being more suitable for practice in the developed world compared to the developing

world (53.9%).

Table 1: Knowledge and attitude of respondents to EBM

Suggestions about EBM	Agreed	Undecided	Dis-agreed	Total
Current research findings are useful in the day-to-day management of my patients	65 (100.0)	0 (0.0)	0 (0.0)	65
Adoption of evidence-based practice places too many demands on my workload	23 (35.4)	4 (6.2)	38 (56.4)	65
EBM is patient centered	56 (86.1)	4 (6.2)	5 (7.7)	65
EBM is of limited value in paediatric practice	3 (4.6)	4 (6.2)	58 (89.3)	65
Literature and research findings are useful in paediatric practice	62 (95.4)	3 (4.6)	0 (0.0)	65
I need to increase the use of evidence in my daily practice	50 (76.9)	15 (23.1)	0 (0.0)	65
I am interested in learning skills to incorporate EBM in my practice	60 (92.3)	5 (7.7)	0 (0.0)	65
EBM improves the quality of patient care	58 (89.3)	6 (9.2)	1 (1.5)	65
EBM does not take into consideration the limitations in clinical practice	36 (55.4)	7 (10.8)	22 (33.8)	65
Reimbursement rate will increase with incorporation of EBM into paediatric practice	12 (18.5)	40 (61.4)	13 (19.9)	65
Strong evidence is lacking in most interventions used	23 (35.5)	15 (23.0)	27 (41.5)	65
EBM should form the basis of decision-making in clinical practice	61 (93.8)	2 (3.1)	2 (3.1)	65
EBM does not take into consideration, patients' preferences	25 (38.5)	7 (10.8)	33 (50.7)	65
EBM is useful in designing the medical curriculum	54 (83.1)	11 (16.9)	0 (0.0)	65
EBM is most useful for paediatric practice in the developed world	35 (53.9)	11 (16.9)	19 (29.2)	65

EBM – Evidence-based Medicine

In the assessment of the skills required to use EBM, Table 2 shows that some of the respondents did not understand but will like to learn heterogeneity (44.6%) while 46.1%, 52.3%, 50.8%, 46.1%, 50.8% and 64.6% understood meta-analysis, systematic review, assessment of publication bias, literature search strategies, study designs and clinical significance of study findings respectively. The highest proportion of the respondents demonstrated the willingness to teach literature search strategies (41.5%) while only 13.8% understood and could teach the use of Cochrane Library Database.

In Table 3 lack of the skills for understanding statistics and lack of skills for locating best research evidence were the leading perceived barriers to the use of EBM as identified by 81.6% and 73.8% respectively of the respondents. The fear of medicolegal tussle arising from the use of EBM was identified by only 27.7% as a barrier to the use of EBM. The proportions of respondents who agreed or disagreed with lack of time and lack of conclusive evidence as barriers to the use of EBM were almost comparable (44.6%). All the respondents agreed

that EBM should be incorporated into Continuing Medical Education activities and into postgraduate medical curriculum while only 56 (86.2%) agreed that EBM should be incorporated into medical undergraduate training curriculum whereas 9 (13.8%) were neutral.

Table 2: Assessment of the skills required for the use of EBM among the respondents

	It is not important I understand	I do not understand but will like to	I have some understanding	I understand and could teach others
Relative Risk	0 (0.0)	23 (35.4)	23 (35.4)	19 (29.3)
Meta-analysis	0 (0.0)	21 (32.3)	30 (46.1)	14 (21.5)
Systematic Review	0 (0.0)	15 (23.1)	34 (52.3)	16 (24.6)
Heterogeneity	0 (0.0)	29 (44.6)	20 (30.8)	16 (24.6)
Publication bias	0 (0.0)	20 (30.8)	33 (50.8)	12 (18.5)
Literature Search Strategies	0 (0.0)	8 (2.3)	30 (46.1)	27 (41.5)
Study designs	0 (0.0)	10 (15.4)	33 (50.8)	22 (33.8)
Evaluating the validity of a study	0 (0.0)	21 (32.3)	26 (40.0)	18 (27.7)
The clinical significance of study results	0 (0.0)	7 (10.8)	42 (64.6)	16 (24.6)
Using the Cochrane Library Database	0 (0.0)	25 (38.5)	31 (47.7)	9 (13.8)

EBM – Evidence-based Medicine

Table 3: Perceived barriers to the use of EBM

	Agreed	Undecided	Disagreed
Lack of time	29 (44.6)	6 (9.2)	30 (46.2)
Lack of conclusive evidence	29 (44.6)	8 (12.3)	28 (43.1)
Lack of computing resources	37 (56.9)	5 (7.7)	23 (35.4)
Lack of access to electronic databases	37 (57.0)	8 (12.2)	20 (30.8)
Lack of skills for locating best research evidence	48 (73.8)	5 (7.7)	12 (18.5)
Lack of skills for understanding statistics	53 (81.6)	0 (0.0)	12 (18.4)
Lack of incentive for using EBM	22 (64.7)	8 (12.3)	15 (23.0)
Fear of medico-legal tussles arising from practice	18 (27.7)	11 (16.9)	36 (55.4)

EBM – Evidence-based Medicine

Assessment of knowledge of the respondents

The knowledge scores for the entire study population ranged between 64.0% and 88.0% with overall mean score of $77.6\% \pm 5.6\%$. The mean scores for the various groups of participants were comparable as shown in Table 4. The assessment of knowledge showed that 52 (80.0%) and 13 (20.0%) participants had high and low knowledge scores respectively. Table 5 shows that the proportions of respondents with high knowledge score were comparable across groups: duration of practice ($p = 0.139$), sector of practice ($p = 0.936$) and setting of practice ($p = 0.710$). In addition, EBM use in practice and prior EBM training were also not significantly associated with high knowledge scores ($p = 0.335$ and $p = 0.947$ respectively).

Table 4: Comparison of mean knowledge scores among comparison groups

Parameters	Groups	Mean scores (%)	t	p-values
Duration of practice	Group I	79.0 ± 4.5	1.583	0.118
	Group II	76.7 ± 6.1		
Sector of practice	Public	78.1 ± 5.4	1.347	0.183
	Private	75.6 ± 0.4		
Setting of practice	Academic	76.5 ± 4.5	-1.604	0.114
	Non-academic	76.0 ± 1.4		
EBM Use in clinical practice	Yes	77.8 ± 5.8	0.734	0.466
	No	76.0 ± 1.4		
EBM Training	Yes	78.6 ± 4.4	1.087	0.281
	No	77.0 ± 6.2		

Table 5: Relationship between the professional characteristics of the respondents and their knowledge of EBM

Characteristics	High Score	Low Score	Statistics
Duration of practice	Group I (n = 24)	22 (91.7)	2 = 2.184; p = 0.139*
	Group II (n = 41)	30 (73.2)	
Sector of practice	Private (n = 12)	9 (75.0)	2 = 0.006; p = 0.936*
	Public (n = 53)	43 (81.1)	
Setting of practice	Academic (n = 33)	27 (81.8)	2 = 0.138; p = 0.710
	Non-academic (n = 32)	25 (78.1)	
EBM Use in practice	No (n = 6)	6 (100.0)	FE = 1.562; p = 0.335
	Yes (n = 59)	46 (78.0)	
EBM Training	No (n = 41)	32 (78.0)	2 = 0.037; p = 0.847*
	Yes (n = 24)	20 (83.3)	

*Yate's Correction applied; FE = Fisher's Exact Test
EBM – Evidence-based Medicine

Relationship between perceived barriers to EBM and professional characteristics of the respondents

In Table 6a, the proportions of respondents who agreed that lack of time was a barrier to EBM were comparable across groups. Significantly higher proportions of Group I respondents and respondents who were trained in EBM disagreed that lack of conclusive evidence was a barrier to EBM ($p = 0.025$ and $p = 0.004$ respectively). Higher proportions of respondents who worked in non-academic settings and those who were trained in EBM disagreed about lack of computing resources being a barrier to EBM ($p = 0.045$ and $p = 0.023$ respectively). Higher proportions of respondents in Group I, those in non-academic settings and those who were trained on EBM also disagreed that lack of access to electronic

database was a barrier to EBM use ($p = 0.007$, $p = 0.002$ and $p < 0.001$ respectively).

Table 6a: Relationship between professional characteristics of the respondents and the perceived barriers to the use of EBM in Paediatric Practice

<i>Lack of time</i>		<i>Agreed</i>	<i>Undecided</i>	<i>Disagreed</i>	<i>Statistics</i>
Duration	Group I(n = 24)	12 (50.0)	0 (0.0)	12 (50.0)	FE = 3.819; P = 0.155
	Group II (n = 41)	17 (41.5)	6 (14.6)	18 (43.9)	
Setting	Academic (n= 33)	15 (45.5)	2 (6.0)	16 (48.5)	2 = 0.664; P = 0.766
	Non-Academic (n = 32)	14 (43.8)	4 (12.5)	14 (43.8)	
EBM Trained	No (n = 41)	19 (46.3)	3 (7.4)	19 (46.3)	2 = 0.561; P = 0.804
	Yes (n = 24)	10 (41.7)	3 (12.5)	11 (45.8)	
<i>Lack of conclusive evidence</i>					
Duration	Group I(n = 24)	10 (41.7)	0 (0.0)	14 (58.3)	FE = 7.041; P = 0.025
	Group II (n = 41)	19 (46.3)	8 (19.6)	14 (34.1)	
Setting	Academic (n= 33)	14 (42.4)	6 (18.2)	13 (39.4)	2 = 2.162; P = 0.376
	Non-Academic (n = 32)	15 (46.9)	2 (6.2)	15 (46.9)	
EBM Trained	No (n = 41)	21 (51.2)	8 (0.0)	12 (29.3)	FE = 10.675; P = 0.004
	Yes (n = 24)	8 (33.3)	0 (0.0)	16 (66.7)	
<i>Lack of computing resources</i>					
Duration	Group I (n = 24)	12 (50.0)	0 (0.0)	12 (50.0)	FE = 5.190; P = 0.06
	Group II (n = 41)	25 (61.0)	5 (1.2)	11 (26.8)	
Setting	Academic (n= 33)	19 (57.6)	5 (15.2)	9 (27.2)	FE = 5.958; P = 0.045
	Non-Academic (n = 32)	18 (56.2)	0 (0.0)	14 (43.8)	
EBM Trained	No (n = 41)	26 (63.4)	5 (12.2)	10 (24.4)	FE = 7.071; P = 0.023
	Yes (n = 24)	11 (45.8)	0 (0.0)	13 (54.2)	
<i>Lack of access to electronic database</i>					
Duration	Group I(n = 24)	12 (50.0)	0 (0.0)	12 (50.0)	FE = 9.612; P = 0.007
	Group II (n = 41)	25 (61.0)	8 (19.5)	8 (19.5)	
Setting	Academic (n= 33)	19 (57.6)	8 (24.2)	6 (18.2)	FE = 11.725; P = 0.002
	Non-Academic (n = 32)	18 (56.3)	0 (0.0)	14 (43.7)	
EBM Trained	No (n = 41)	28 (68.3)	8 (19.5)	5 (12.2)	FE = 19.198; P < 0.001
	Yes (n = 24)	9 (37.5)	0 (0.0)	15 (62.5)	

*FE = Fisher's Exact Test; EBM – Evidence-based Medicine

Table 6b: Relationship between professional characteristics of the respondents and the perceived barriers to the use of EBM in Paediatric Practice

<i>Lack of skill to locate best research evidence</i>		<i>Agreed</i>	<i>Undecided</i>	<i>Disagreed</i>	<i>Statistics</i>
Duration	Group I(n = 24)	14 (58.3)	0 (0.0)	10 (41.7)	FE = 14.312; P < 0.001
	Group II (n = 41)	34 (82.9)	5 (12.2)	2 (4.9)	
Setting	Academic (n= 33)	28 (84.8)	3 (9.1)	2 (6.1)	2 = 6.853; P = 0.033
	Non-Academic (n = 32)	20 (62.5)	2 (6.3)	10 (31.2)	
EBM Trained	No (n = 41)	34 (82.9)	5 (12.2)	2 (4.9)	FE = 14.312; P < 0.001
	Yes (n = 24)	14 (58.3)	0 (0.0)	10 (41.7)	
<i>Lack of skills for understanding statistics</i>					
Duration	Group I(n = 24)	16 (66.7)	0 (0.0)	8 (43.3)	2 = 4.134*; P = 0.042
	Group II (n = 41)	37 (90.2)	0 (0.0)	4 (9.8)	
Setting	Academic (n= 33)	21 (63.6)	2 (6.1)	10 (30.3)	2 = 5.277*; P = 0.022
	Non-Academic (n = 32)	22 (68.7)	8 (25.0)	2 (6.3)	
EBM Trained	No (n = 41)	37 (90.2)	0 (0.0)	4 (9.8)	2 = 4.134*; P = 0.042
	Yes (n = 24)	16 (66.7)	0 (0.0)	8 (33.3)	
<i>Lack of incentives for using EBM in practice</i>					
Duration	Group I(n = 24)	20 (83.3)	0 (0.0)	4 (16.7)	FE = 7.567; P = 0.02
	Group II (n = 41)	22 (53.7)	8 (19.5)	11 (26.8)	
Setting	Academic (n= 33)	22 (66.7)	4 (12.1)	7 (21.2)	2 = 0.147; P = 0.929
	Non-Academic (n = 32)	20 (62.5)	4 (12.5)	8 (25.0)	
EBM Trained	No (n = 41)	24 (58.5)	8 (19.5)	9 (22.0)	FE = 5.715; P = 0.06
	Yes (n = 24)	18 (75.0)	0 (0.0)	6 (25.0)	
<i>Fear of medicolegal tussles following EBM use</i>					
Duration	Group I(n = 24)	8 (33.3)	0 (0.0)	16 (66.7)	FE = 8.791; P = 0.013
	Group II (n = 41)	10 (24.4)	11 (26.8)	20 (48.8)	
Setting	Academic (n= 33)	10 (30.3)	8 (24.2)	15 (45.5)	2 = 3.395; P = 0.167
	Non-Academic (n = 32)	8 (25.0)	3 (9.3)	21 (65.7)	
EBM Trained	No (n = 41)	13 (31.7)	11 (26.8)	17 (41.5)	FE = 11.727; P = 0.003
	Yes (n = 24)	5 (20.8)	0 (0.0)	19 (79.2)	

*Chi-Square with Yate's Correction; FE = Fisher's Exact Test; EBM – Evidence-based Medicine

Table 6b shows that higher proportions of Group I respondents, respondents in non-academic settings and respondents trained in EBM disagreed that lack of skills required to locate best research evidence is a barrier to EBM use ($p < 0.001$, $p = 0.033$ and $p < 0.001$ respectively). Lack of skills for understanding statistics was not perceived as a barrier to EBM by significantly higher proportions of Group I respondents, respondents in the academic setting and respondents who were trained on EBM ($p = 0.042$, $p = 0.022$ and $p = 0.042$ respectively). In addition, a significantly higher proportion of Group I respondents agreed that lack of incentives for using EBM in practice was a barrier to the use of EBM ($p = 0.02$). Significantly higher proportions of respondents in Group I and those who were trained in EBM disagreed that there may be medicolegal tussles following EBM use ($p = 0.013$ and $p = 0.003$ respectively).

Discussion

The present study revealed that almost all the respondents (90.8%) used EBM in their clinical practices suggesting a high level of acceptability of EBM in the population. This contrasts with 40% EBM use rate reported among doctors in Jordan⁹ and 5% each reported in Sudan and Jordan.^{7,10} The observed difference may be explained in terms of differences in the degree of exposure to EBM during professional workshops and conferences over time. However, the high use rate of EBM in the present study contrasted sharply with the low proportion of participants (36.9%) who had received training on EBM. This pattern was similar to the findings among a cohort of Nigerian specialist trainees, where 96.6% were familiar with EBM but only 38.8% had been formally trained on EBM.¹¹ This low rate of training was similar to 15% and 24% previously reported in Sudan and Sri Lanka respectively.^{10,12} This observation may be related to the poor awareness of training opportunities as currently provided by organisations such as the various national and regional Cochrane Centres. It is plausible that paucity of funds and logistic supports may actually limit the number of people that such organisations could admit for training at a time. Interestingly, the higher cadre participants in the present study were four times more likely to have received training on EBM compared to the junior cadre participants. The reason for this observation is obscure but it is plausible that the senior participants had better access to training opportunities by virtue of hierarchy in the profession. The universal recommendation of the participants in the present study that training on EBM should be incorporated into postgraduate training curricula may provide a solution to the lopsidedness in the pattern of EBM training in relation to the duration of practice.

Fourth-fifth of the participants in the present study had high knowledge of what EBM entails and what purpose it serves. This is not surprising as the overall mean knowledge score was 77.6% compared to 24% recorded

in a population of Iranian doctors.¹³ Although the relatively high knowledge score observed in the present study is reassuring, it may not be a perfect proof of deep knowledge of EBM, given the low rate of training reported in the same population. Almost all the participants (92.3%) were willing to learn the skills required to incorporate EBM into their routine clinical practice and this may be a reflection of the acceptability of EBM in clinical practice.

The major perceived barriers to the use of EBM in the present study included the lack of skills for understanding statistics, lack of skills for locating best research evidence and lack of incentives for using EBM. Interestingly, the fear of medicolegal tussles arising from the practice of EBM was the least identified barrier. The lack of time was not perceived a leading barrier in the present study unlike other previous studies in Jordan,⁹ Sudan,¹⁰ Saudi Arabia,¹⁴ Norway,¹⁵ and Sri Lanka.¹² This may reflect the heterogeneous nature of the participants in the various cited studies. Lower cadre doctors are likely to be busier in terms of clinical duties than the higher cadre doctors, who are mostly involved in administrative duties alongside academics with some involvement in clinical duties. Therefore, studies focused on higher cadre doctors, like the present study, are not likely to report lack of time or overwhelming workload as perceived barriers to the practice of EBM. However, the challenge of lack of time, though coming a distant sixth position on a list of eight in the present study, may be addressed by incorporating research day-off on a weekly or bi-weekly basis into the duty schedule for doctors at all levels.

Lack of understanding of statistics and lack of skills for searching and locating best research evidence are training issues which frequent Continuing Medical Programmes can address to a large extent. Access to internet EBM databases such as the Cochrane Library and the World Health Organisation Reproductive Health Library, may need to be improved, just as an earlier study in Bosnia Herzegovina¹⁶ reported that 34.6% of doctors did not know how to use the Cochrane database similar to 38.5% observed in the present study. Institutions should be encouraged to subscribe to these databases and facilitate easy and hitch-free access and use by doctors, particularly when they are not kept busy with clinical duties. Lack of evidence as a perceived barrier to the use of EBM had been reported by Al-Almaie in Saudi Arabia.¹⁷ That may be related to the fact that the Cochrane Database of Systematic Reviews display many systemic reviews with inconclusive findings arising from either small number of study participants or poor design of the primary randomized controlled trials. Therefore, this dearth of studies with conclusive findings should not deter doctors from the use of the available facts. In addition, this should also stimulate more research in the form of well-designed, high quality randomized controlled trials to answer some of the yet unanswered research questions.

Al-Omari⁷ reported threat to clinical freedom as a perceived barrier to the use of EBM among doctors in Jordan but that fear of medicolegal tussle surrounding the use of EBM did not appear strong in the present study. This difference may be related to the relative rate of medical litigations in each environment. Where treatment protocols have been drawn for routine clinical practice from the best quality evidences available, it becomes difficult to defend any other clinical decision aside the recommendations of the treatment protocol and that may predispose to litigations. In Nigeria, the process of generating treatment protocols for paediatric practice are still on-going, hence there is no pressure for mandatory adherence to specific practices yet. Therefore, medicolegal issues may not yet be perceived as a threat to the use of EBM.

The Group I participants (relatively higher in the hierarchy) and those who had received trainings on EBM were more likely to disagree with the perceptions of lack of computing skills, lack of access to electronic databases, lack of skill to locate the best research evidence, lack of skill for understanding statistics, lack of incentives and fear of medicolegal tussles as barriers to EBM. This observation may be explained in terms of the role of experience on perceptions. The perceived barriers are

usually routinely addressed during EBM training and with practice, every researcher learns that the perceived barriers can be adequately overcome.

Conclusion

In conclusion, the knowledge of this population of Nigerian paediatricians about EBM was good but most of the participants crave better understanding of the statistical aspects of systematic reviews and EBM. The participants also recommended the inclusion of EBM training in both the undergraduate and postgraduate medical curricula as well as in Continuing Medical Education programmes. With adequate training, the perceived barriers to use of EBM will be addressed and appropriate solutions will be provided.

The electronic method of gathering the responses of the participants is acknowledged as a limitation to the study as it may explain the low response rate.

Conflict of interest: None

Funding: None

References

1. Oladapo OT. Research synthesis: Implications for clinical practice in Nigeria. *International J Multi disciplinary Research 2012*; *V(1&2)*: 4-6.
2. Rosenberg WM, Sackett DL. On the need for evidence based medicine. *Therapie 1996*; *51(3)*: 212-217.
3. Collins JA. Clinical research evidence and clinical practice. *Hum Reprod 1997*; *12(9)*: 1847-1850.
4. Mahmoud AO, Ayanniyi AA, Lawal A, Omolase CO, Ologunsua Y, Samaila E. Survey of medical specialists on their attitudes to and resources for health research in Nigeria. *Ann Afr Med 2011*; *10(2)*: 144-149.
5. Mahmoud AO, Ayanniyi AA, Lawal A, Omolase CO, Ologunsua Y, Samaila E. Perceptions of Nigerian medical specialists on research. *J Public Health Afr 2011*; *2(1)*: e1. Doi: 10.4081/jphia.2011.e1
6. Sackett DL, Rosenberg WM. Th need for evidence-based medicine. *J R Soc Med 1995*; *88(11)*: 620-624.
7. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. Cochrane Book Series. Wiley-Blackwell. Chichester, West Sussex, England. 2008.
8. Barghouti F, Halaseh L, Said T, Mousa AH, Dabdoub A. Evidence-based medicine among Jordanian family physicians: awareness, attitude and knowledge. *Can Fam Physician 2009*; *55(7)*: e6-e13.
9. Zeidan AZ, Behairy MM. Knowledge, attitude and practice of Evidence-based medicine among residence doctors in Sudan. *Sudan J Med Sci 2010*; *5(3)*: 2007-2012.
10. Al Omari M, Khader Y, Jadalalah K, Dauod AS, Al-Shdifat AA, Khasawneh NM. Evidence-based medicine among hospital doctors in Jordan: awareness, attitude and practice. *J Eval Clin Pract 2009*; *15(6)*: 1137-1141.
11. Fawole A, Oladapo O, Enahoro F, Akande E. Acceptance of evidence-based reproductive health care among postgraduate specialist trainees in Nigeria. *Int J GynaecolObstet 2008*; *102(1)*: 3-7.
12. Abeysena C, Jayawardana P, Wickremasinghe R, Wickramasinghe U. Evidence-based medicine knowledge, attitudes and practices among doctors in Sri Lanka. *J Evid Based Med 2010*; *3(2)*: 83-87.
13. Rashidbeygi M, Sayehmiri K. knowledge and attitude of physicianstowarss evidence based medicine in Ilam, Iran. *Iran Red Crescent J 2013*; *15(9)*: 798-803.
14. Al-Musa HM. Knowledge, perceptions, attitude and educational needs of physicians to evidence based medicine in South-Western Saudi Arabia. *Saudi Med J 2010*; *31(3)*: 308-312.
15. Ulvenes LV, Aasland O, Nylenna M, Kristianseh IS. Norwegian physicians' knowledge of and opinions about evidence-based medicine: cross-sectional study. *PLoS One 2009*; *4(11)*: e7828. Doi: 10.1371/journal.pone.0007828.

16. Mahmi -KaknjoM, Kadi D, Hodžić H, Spahi -Sarajli S, Hadžić E, Ademovi E. Awareness, knowledge, use and attitude towards evidence based medicine in a developing country: Survey of physicians in a canton in Bosnia and Herzegovina. *Croat Med J* 2015; 56(6): 558-566.
17. Al-Almaie SM, Al-Baghli N. Barriers facing physicians physicians practicing evidence based medicine in Saudi Arabia. *J ContinEduc Health Prof* 2004; 24(3): 163-170.