



NIGERIAN JOURNAL OF PAEDIATRICS

VOLUME 53 | NUMBER 3 | JULY - SEPTEMBER 2026

<https://www.njpaediatrics.com>

PRINT: ISSN 0302-4660

ONLINE: ISSN 2814-2985

SPECIAL ARTICLE Transforming Nigeria's Health System to Address Contextual Challenges in Child Healthcare Training and Practice
Esangbedo Dorothy O

REVIEWS A Narrative Review on the Diagnostic Features in Sickle Cell Nephropathy with a Focus on Urine Protein-Creatinine Ratio

Akinsete Adeseye M, Fakile Ugonna O, Ezeaka Nonye, West Timipah, Mbanefo Ngozi, Saleh Ahmed F, Ojika Blessing, Alakaloko Adebimpe, Damola-Okesiji Adebola, Temiye Edamisan

A Systematic Review of Superficial Fungal Infections in Nigerian Children and a Case for Inclusion in Neglected Tropical Diseases
Makarfi Hauwa'u U, Mohammed Tahir T

ORIGINAL ARTICLE A Comparative Study of Socio-demographic Characteristics and Risk Factors of Epilepsy Between Children with and without Co-existing Cerebral Palsy in Sokoto, Northwest Nigeria

Mikailu Abubakar J, Ahmed Hamidu, Ahmad Murtala M, Jiya Fatima B, Yusuf Abduganiyyu, Waziri Usman, Ahmed Hadiza K, Ishaq Fatima A, Ibrahim Suleiman M, Surajo Ibrahim

Awareness, Knowledge and Attitude of Medical Students to Room Air Use in Neonatal Resuscitation at the Igbinedion University, Okada, Nigeria

Ibhanesebhor Samuel E, Umeh Evan I

Enhancing Immunisation Communication: Perspectives from Doctors and Vaccinators in Kano, Nigeria

Farouk Zubaida L, Ibrahim Hafsat U, Kabir Halima, Ali Aminu, Garba Maria A, Tabansi Petronila, Tagbo Beckie, Ogunrinde Olufemi G

Apolipoprotein B-100/Apolipoprotein A-I Ratio in overweight adolescents with or without metabolic syndrome

Barretto Neto Augusto C, Silva Vivianne SD, Lima Vera L, Alcides da Silva D

Comparative Assessment of the Nutritional Status of Institutionalised and Non-Institutionalised Nigerian Children Using

Anthropometric Parameters

Oji-Onuoha Stella N, Ughasoro Maduka D, Izuka Mildred O, Chukwudi Ndubuisi K, Oji Onuoha, Jibrin Alkasim M

Caregivers' Voices: Experiences and Satisfaction with Paediatric Care in a Nigerian Tertiary Hospital

Mustapha Samaha S, Musa Aishatu Z, Musa Sulaiman A, Lawal Nurat O, Avong Olusola, Shuaibu Kaltimi

Association Between Sleep Quality and Overweight/Obesity Among Secondary School Adolescents in Yenagoa Local Government

Area, Bayelsa State, Nigeria

Onyeka Chiemerie A, Okosun Ofure A, Akinbami Felix O

The Use of Middle Finger Length as a Predictor of Standing Height Among Adolescents in Yenagoa, Bayelsa State, Nigeria

Okumoko Maureen A, Binafeigha Ihekerenna J, Akinbami Felix O

Point of Admission Hypothermia and Its Correlates Among Hospitalized Neonates at the Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria

Abdulgafar Habeebllahi A, Adeyemo Ibrahim O, Adamu Asman, Onankpa Ben, Yusuf Tahir, Akpelu Echieme, Atusu Yasmine, Isezuo Khadijat

Social Determinants of Health Among Children and Adolescents with Sickle Cell Disease Attending Two Tertiary Hospitals in Southwest Nigeria

Olasinde Yetunde T, Adetoye Mayowa M, Olasinde Adeola, Afolayan Odunayo O, Agelebe Efeturi, Onigbinde Michael O, Adeola Samuel A, Adeagbo Victor O, Tijani Oluwadamilola E, Idowu Ajibola

Child Adoption Practices: Emerging Child Welfare Concerns in a Nigerian Community

Odita Amalachukwu O, Mbachu Chioma NP, Ezeasomba Fortress I, Anioto Ezinne C

Paediatric and Adolescent Breast Ultrasound in Ibadan, Nigeria: Imaging Findings and Correlation of BI-RADS with Histopathology

Adeniji-Sofoluwe Adenike T, Onyema Adaora, Oni Oluwatomi Y, Obajimi Millicent O

Association of Neck Circumference with Pulse Pressure Among School-aged Children in Ibadan, Nigeria: A Cross-Sectional Study
Orimadegun Adebola E, Okhuaihesuyi Osaruese

CASE REPORT Severe Hypertriglyceridaemia with Acute Kidney Injury in Paediatric Diabetic Ketoacidosis: A Case Report

Ahmad Raji M, Nwalorzie Chinomso, Ononiwu Uche, Oguzie Chioma

Neonatal Diabetic Ketoacidosis Masquerading as Sepsis in a Nigerian Infant: A Case Report

Solomon Amos, Omachi Adah P, Eze Edith C, Abdullahi Usman, Mammud Abubakar, Abubakar Usman

Experience With Intra-Lesional MMR Vaccine in the Treatment of Anogenital Warts in Children: A Report of Three Cases

Lamina Amotunur B, Odusote Olatunde A, Ubuane Peter O

EDUCATIONAL SERIES CLINICAL QUIZ
Senbanjo Idowu O.

OFFICIAL JOURNAL OF THE PAEDIATRIC ASSOCIATION OF NIGERIA



Nigerian Journal of Paediatrics 2026 (September); Volume 53(3).

<https://dx.doi.org/10.63270/njp.v53i3.2000072>.

A Systematic Review of Superficial Fungal Infections in Nigerian Children and a Case for Recognition as Neglected Tropical Diseases

Makarfi Hauwa'u U¹, Mohammed Tahir T²

¹Department of Paediatrics, College of Medical Sciences, Ahmadu Bello University, Zaria, Nigeria.

²Department of Medicine, College of Medical Sciences, Ahmadu Bello University, Zaria, Nigeria.

Correspondence

Dr Makarfi Hauwa'u U. Department of Paediatrics, College of Medical Sciences, Ahmadu Bello University, Zaria, Nigeria. E-mail: humakarfi@yahoo.com ; ORCID – <https://orcid.org/0000-0003-4624-6369>.

Abstract

Background: Superficial fungal infections (SFIs) are among the most common infectious dermatoses in children worldwide. In low- and middle-income countries such as Nigeria, they remain highly prevalent, yet under-prioritised in public health policy.

Objective: To systematically review the epidemiology, clinical spectrum, risk factors, and public health implications of superficial fungal infections among Nigerian children and assess whether they meet criteria consistent with neglected tropical diseases (NTDs).

Methods: We conducted a systematic search of PubMed, African Journals Online, Google Scholar, and institutional repositories for studies published from 2005 to 2025 that reported primary epidemiological data on SFIs in Nigerian children. Inclusion criteria comprised studies reporting prevalence and clinical patterns in individuals ≤ 18 years, and available in English. Data were extracted on study location, population studied, sample size, prevalence rates and clinical types of SFI, common organisms and risk factors.

Results: Thirty-three studies met the inclusion criteria. Reported prevalence ranged from 4.7% to 75.7%, with most studies documenting rates of 20–60%. Tinea capitis was consistently the predominant infection, particularly among children aged 6–12 years, with male predominance. Risk factors included overcrowding, poor hygiene, low socioeconomic status, and sharing of personal grooming tools. Diagnostic heterogeneity and limited mycological confirmation were common limitations. The review showed some similarities between SFIs and the officially recognised neglected tropical diseases.

Conclusion: Superficial fungal infections are highly prevalent among Nigerian children but remain under-recognised in national disease prioritisation frameworks. There's a need to designate these infections as neglected tropical diseases in order to improve their control and prevention.

Keywords: *Candida*, *Dermatophytosis*, *Low socioeconomic status*, *Malassezia*, *Superficial mycosis*, *Tinea capitis*.

Introduction

Superficial fungal infections (SFIs) are caused by dermatophytes — fungi that invade keratinised tissues (skin, hair, nails) — and non-dermatophyte organisms such as *Candida* and *Malassezia*.¹ Despite being a recognised global public health problem, most

symptomatic patients never seek care.^{2,3} SFIs can be disfiguring and cause intractable itch, significantly reducing quality of life.² SFIs occur across all demographics but disproportionately affect children. A systematic review across Africa recorded an overall SFI prevalence of 17%, with Nigerian children

averaging over 23%.⁴ The contributing factors include low socioeconomic status, overcrowding, and animal contact, compounded by psychosocial stigma. Yet childhood dermatophytosis has attracted little targeted public health intervention compared to conditions like scabies and leprosy — despite sharing key features with neglected tropical diseases (NTDs).

The World Health Organisation (WHO) defines NTDs as a diverse group of transmissible infections affecting over 1.5 billion impoverished people lacking access to clean water, sanitation, and quality healthcare.⁵ Currently, only certain deep mycoses are classified as NTDs; no dermatophytoses are yet included.⁶ NTD-defining characteristics include causation of more disability than mortality, chronic clinical course, longstanding existence with minimal research, tropical/subtropical distribution, and strong association with poverty⁶ - all of these apply equally to SFIs. This review therefore argues that the same intervention strategies used for NTDs should be applied to childhood SFIs to improve quality of life for affected children and communities.

This systematic review aims to examine the available literature from Nigeria on childhood

SFIs and describe the burden and determinants to make a case for consideration of childhood dermatophytosis as a possible NTD.

Review Methods

This systematic review examined published literature on SFIs in Nigerian children retrieved from PubMed, African Journals Online (AJOL), Google Scholar, and institutional repositories. Search terms included combinations of "superficial fungal infections", "tinea capitis", "dermatophytosis", "pityriasis versicolor", "Nigeria", "children", and "school", restricted to studies published between 2005 and December 2025.

Inclusion criteria: Nigerian studies with participants under 18 years, reporting primary prevalence or clinical pattern data, and available in English.

Exclusion criteria: Adult-only or mixed adult-child populations, non-primary data sources (reviews, case reports, conference abstracts), invasive/systemic mycoses only, or laboratory data without clinical correlation.

The extracted variables included the author, year, location, population, sample size, participant age, prevalence rate, most common infection, predominant organism, and reported risk factors. A representation of the search schema is shown in Figure 1.



Figure 1: PRISMA flow diagram for the search criteria.

Results

A total of 33 full-text articles were included in the qualitative analysis; the majority of these studied primary school children. The combined sample includes >100,000 children from urban, rural, agrarian, nomadic, and commercial communities. Meta-analysis (quantitative analysis) was not performed due to the high heterogeneity in the study methods and reporting of clinical patterns and risk factors. Table I summarises the characteristics of the included studies.

Epidemiology and prevalence of childhood SFIs

Across the 33 studies from several regions of Nigeria, the prevalence of SFIs among children was high and variable, ranging from 4.7% to 75.7%, depending on diagnostic methods (clinical versus culture), region, and socio-economic context. Most studies reported prevalence rates from 20% to 60%, with nine studies below 20% and six above 60%. Studies were more concentrated in the south-east and south-west regions, with no studies from the north-east; however, one or more studies from every other region reported prevalence within the 20–60% range.^{7,13,15,25,38} All the reviewed studies employed a cross-sectional design, predominantly recruiting primary school children. Notable exceptions include Nweze *et al.*,¹⁷ who studied children in nomadic herdsman camps in south-east Nigeria (prevalence 41%), and hospital-based studies by Okafor *et al.* and Altraide and Alex-Hart, which reported lower prevalences of 6.7% and 8.1%, respectively.^{20,31} One case-control study for dermatophytosis originally reported 91.35%, which was recalculated as 45.1% to account for the study design and is the value used in this review.²⁹

Despite methodological heterogeneity, the available data consistently indicate that SFIs are highly prevalent among Nigerian children, with most prevalence exceeding 20% across all sub-regions. Regarding age distribution, most studies reported the highest burden among

primary school-aged children (6 to 12 years), with fewer infections recorded above 12 years of age.^{14,36} A minority of studies found no significant age-related differences,^{19,21,25,31} while others identified peak infection in older (10 to 14 years) or pre-school (below 6 years) age groups.^{7,15,38} Male sex was the predominant risk demographic in most studies, with only a few reporting no significant gender difference.^{7,17,25,31} Collectively, the evidence supports a predilection of SFIs for male children aged 6 to 12 years.

Clinical spectrum of SFIs in Nigerian children

Tinea capitis emerged consistently as the predominant SFI across reviewed studies, with a wide prevalence range of 3.9% to 85.4%. Only one study identified tinea corporis as the leading type (50%), with tinea capitis next at 23.3%.³¹ Non-inflammatory forms of dermatophytosis accounted for more than 95% of documented cases of tinea capitis, though the relative proportion of grey patch versus black dot types varied across studies. Inflammatory presentations, such as kerion, were uncommon (1% to 2%).^{23,36} Other SFIs occurred at considerably lower frequencies: tinea corporis (0.6% to 8.5%), tinea faciei (0.5% to 1.1%), tinea pedis and unguium (0.1% to 0.9%), and pityriasis versicolor (2.4% to 6.1%).^{10,13,14,23,31,36}

Aetiologic agents (Dermatophytes) of SFIs

More than three-quarters of the reviewed studies successfully isolated causative fungal organisms. Three dermatophyte genera predominated: *Trichophyton spp.*, *Microsporum spp.*, and *Epidermophyton spp.* The most commonly reported leading isolates were *T. mentagrophytes* (five studies),^{7,17,22,30,39} *T. rubrum* (five studies),^{21,25,26,29,33} *M. audouinii* (three studies),^{16,18,23} *T. tonsurans* and *T. soudanense* (two studies each),^{14,15,35,36} and *M. canis*, *M. gypseum*, and *E. floccosum* (one study each).^{12,27,37}

Table 1: Characteristics of reviewed studies

Authors / Year	Location	Population	Sample size	Age Range	Prevalence	Most Common Infection	Most Common Dermatophytes	Major Risk Factors
Anosike <i>et al.</i> , 2005, ⁷	Ebonyi	Primary pupils	279	5-13yrs	21.1%	Not documented	<i>T. mentagrophytes</i> 22%	Age group
Enendu & Ibe, 2005 ⁸	Uli, Anambra	Primary school	1,020	4-15yr	13.9%	Tinea capitis 15.5%	<i>T. tonsurans</i> 27.3%	Not documented
Nweze & Okafor, 2005 ⁹	Anambra	Primary schools	1,624	4-16 yrs	5.9%	Tinea capitis 45%	<i>T. tonsurans</i> 37.5%	Age group, rural areas
Ogunbiyi <i>et al.</i> , 2005, ¹⁰	Ibadan	Primary schools	1,066	4-15 yrs	15.2%	Tinea capitis 14.5%	ND	Large family; pets; shared clothing
Uneke <i>et al.</i> , 2005 ¹¹	Ebonyi	Primary and Secondary schools	500	7-17	62.4%	Tinea capitis 51.8%	ND	No statistically significant risk factors
Mbata & Nwajagu, 2006 ¹²	Awka	Nursery and Primary school	2,117	1-13	4.7%	Tinea capitis 3.9%	<i>M. gypsum</i> 33.3%	Not documented
Oyedeji <i>et al.</i> , 2006 ¹³	Ijesha land	Primary school	1,120	6-12	21.1%	Tinea capitis 43.3%	ND	Age group, Male sex
Ngwogu & Otokunefor, 2007 ¹⁴	Abia (rural)	Primary schools	4,287	4-18 yrs	20.4%	Tinea capitis 13.7%	<i>T. soudanense</i> , 26.2%	Not documented
Ayanbimpe <i>et al.</i> , 2008 ¹⁵	Benue & Plateau)	Primary pupils	28,505	3-16 yrs	31.2%	Tinea capitis 31.2%	<i>T. soudanense</i> 30.6%.	Age group, Male sex
Emele & Oyeka, 2008 ¹⁶	Anambra	Primary schools	47,723	2.5-15	9.4%	Tinea capitis 9.4%	<i>M. audouinii</i> 42%	No statistically significant risk factors
Nweze, 2010 ¹⁷	Fulani/Hausa camps Enugu	Nomadic camps	390	4-17	41.5%	Tinea capitis 69.1%	<i>T. mentagrophytes</i> , 29.1%	Age group
Sanuth & Efuntoye, 2010 ¹⁸	Ago-Iwoye, Ogun	Primary school	1,404	7-12	19.5%	Tinea capitis 82.4%	<i>M. audouinii</i> 14%	Soil/animals, poverty
Amoran <i>et al.</i> , 2011, ¹⁹	Sagamu	Primary school	480	5-16	74.1%	Tinea capitis 41.8%	ND	Male sex, previous infection, sharing towels
Okafor <i>et al.</i> , 2011 ²⁰	Ibadan	Hospital based	402	3mo - 12 yrs	6.7%	Tinea capitis 6.7%	ND	Low SES, male sex
Enemuor <i>et al.</i> , 2013 ²¹	Anyigba, Kogi	Primary pupils	2,184	4-15	6.6%	Tinea capitis 85.4%	<i>T. rubrum</i> 30.3%	Male sex, low parental education
Ayanlowo <i>et al.</i> , 2014, ²²	Sagamu, Ogun State	Primary schools	604	2-17	53%	Tinea capitis 15.4%	<i>T. mentagrophytes</i> 51.7%	Low SES
Oke <i>et al.</i> , 2014 ²³	Ile-Ife, Osun	Primary pupils	800	5-16	35%	Tinea capitis 26.9%	<i>M. audouinii</i> 28%	Male sex, tribe
Ogbu <i>et al.</i> , 2015 ²⁴	Awka South	Primary school	870	5-14	40.6%	Tinea capitis 64.9%	ND	No statistically significant risk factors
Dogo <i>et al.</i> , 2016 ²⁵	Kaduna	Primary schools	100	5-15	45%	Tinea capitis 45%	<i>T. rubrum</i> 28.8%	Family size, bed sharing
Ahmed, 2017 ²⁶	Kaduna	Primary school	100	5-15	45%	Tinea capitis 45%	<i>T. rubrum</i> 27.2%	Age group, number of persons living in the house, barbers

A Systematic Review of Superficial Fungal Infections in Nigerian Children and a Case for Recognition as Neglected Tropical Diseases

Afolabi <i>et al.</i> , 2018 ²⁷	Osun	Primary schools (public vs private)	420	4–16	21.7%	Tinea capitis 21.7%	<i>E. floccosum</i> 12% Non-dermatophytes 33%	Poor hygiene, low SES, public schools, sharing combs, sand play
Damen <i>et al.</i> , 2018 ²⁸	Plateau	Primary school	185	5–12	65.4%	Tinea capitis 65.4%	<i>Trichophyton spp.</i> 49.7%	No statistically significant risk factors
Adesiji <i>et al.</i> , 2019 ²⁹	Osogbo (case-control)	Primary schools	230	4–15	45.6%	Tinea capitis (45.6%)	<i>T. rubrum</i> 34%	Poor hygiene, age group
Anya & Sunday, 2019 ³⁰	Aba	Primary school	100	5–13	53%	Not Documented	<i>T. mentagrophytes</i> 43.4%	Not documented
Altraide & Alex-Hart, 2020 ³¹	Port Harcourt	Hospital-based study	347	6–18	8.17%	Tinea capitis 23.3% Tinea corporis 50%	ND	No statistically significant risk factors
Odeigah <i>et al.</i> , 2020 ³²	Ilorin	Primary pupils	602	5–16	29.9%	Not documented	Dermatophytes 16.7% Non-Dermatophytes 51.7%	Age 9–12, overcrowding, poor hygiene
Ayodele <i>et al.</i> , 2021 ³³	N. Central Nigeria	Primary pupils	301	5–14	75.7%	Tinea capitis 75.7%	<i>T. rubrum</i> 68.9%	Large family size
Ezomike <i>et al.</i> , 2021 ³⁴	Enugu	Primary pupils (urban vs rural)	1,662	5–15	45%	Tinea capitis 73.7%	ND	Urban residence, poor hygiene, Male sex
Olarinoye <i>et al.</i> , 2021 ³⁵	Ilorin	Primary schools	204	4–16	73.5%	Tinea capitis 66.1%	<i>T. tonsurans</i> 41.2%	No statistically significant risk factors
Onubogu <i>et al.</i> , 2022 ³⁶	Ukpor & Nnewi (agrarian vs commercial)	Primary pupils	1,562	5–15	28.9%	Tinea capitis 23.5%	<i>T. tonsurans</i> 37.7%	Male sex, public schools, low socio-economic class
Ukhureigbe <i>et al.</i> , 2022 ³⁷	SW Nigeria	Primary schools	102	1–14	73.5%	Tinea capitis 73.5%	<i>M. canis</i> , 28%	No statistically significant risk factors
Okpo <i>et al.</i> , 2024 ³⁸	Calabar	Primary school	779	4–17	25.9%	Tinea capitis 63.9%	<i>Trichophyton spp.</i> 35.6%	No statistically significant risk factors
Abdu <i>et al.</i> , 2025 ³⁹	Kaduna metropolis	Primary school	400	3–14	29.3%	Tinea capitis 29.3%	<i>T. mentagrophytes</i> 10.8%	No statistically significant risk factors

ND – Not Documented

Non-dermatophyte fungi were the predominant isolates in two studies.^{27,32} No consistent regional distribution pattern was identified, though *Trichophyton spp.* dominated across most regions, with the exception of the south-west, where *Microsporum spp.* predominated in half of the studies.

Associated risk factors of reported SFIs

Multiple risk factors were identified across studies, though fewer underwent formal statistical testing. Age 6 to 12 years was the most frequently implicated age

group,^{7,9,13,15,17,26,29,32} and male sex was consistently associated with higher infection rates.^{13,15,19–21,23,34,36} Low socioeconomic status was significantly associated with SFIs in several studies.^{18,20–22,27,36} Additional factors included large household size, bed-sharing, overcrowding,^{10,25,26,32,33} poor personal hygiene (infrequent bathing, not using soap, poor hair care),^{27,29,32,34} sharing of personal items such as combs and towels, attendance at community barbers, and close-contact play.^{10,19,25–27} Contact with soil or animals was documented in a subset of studies,^{10,18,27} as was attendance at

public schools.^{27,36} A prior episode of SFI was identified as a risk factor for recurrence in one study.¹⁹

Antifungal susceptibility studies for SFIs

Only two studies examined antifungal susceptibility patterns, with limited overlap in isolates and agents tested, precluding meaningful comparison. Notably, shared isolates (*T. rubrum*, *T. mentagrophytes*, *M. ferrugineum*) yielded contradictory susceptibility results between the two studies for itraconazole, terbinafine, and griseofulvin.^{33,39} The data are therefore insufficient to support generalisable conclusions regarding antifungal susceptibility for SFIs in Nigeria.

Co-morbidities of SFIs in Nigerian children

Several studies documented concurrent skin conditions, with scabies reported at a prevalence of 1.1% to 10.8%,^{10,13,19,31} a finding of public health relevance given scabies' designation as a Neglected Tropical Disease (NTD) in 2017.⁵ Scarring or patchy alopecia was documented in 15.7% to 38.8% of affected children in two studies,^{22,35} and psychosocial distress was reported in 50% of affected children by Olarinoye *et al.*³⁵ Two studies examining health-seeking behaviour found that more than 90% of affected children did not access appropriate medical care, and up to 70% were self-medicating or using unorthodox treatments.^{19,35}

Discussion

This review demonstrates that SFIs are highly prevalent among Nigerian school children, with tinea capitis dominating the clinical spectrum. Rates vary across regions and methodologies, yet many studies report prevalence approaching or exceeding that of other endemic dermatoses such as scabies. The burden is compounded by significant treatment gaps, with most affected children receiving no care or inappropriate remedies. The absence of nationwide prevalence studies, standardised diagnostic

criteria, and longitudinal data limits accurate national burden estimation.

The key drivers of this high endemicity include low socio-economic status or poverty, overcrowding and poor personal hygiene. These social determinants highlight the need for improved health advocacy with particular emphasis on strengthening the school-based health screening programme, as emphasised by many of the studies. Unfortunately, there was limited local antifungal susceptibility data to inform evidence-based empirical treatment guidelines for SFIs in Nigeria. Although no mortality was attributed to SFIs, they represent a persistent community health concern owing to significant psychosocial sequelae, cosmetic consequences (e.g., alopecia), household transmission risk and potential for educational disruption from constant itching/irritation.

Should SFIs be considered Neglected Tropical Diseases?

SFIs in Nigerian children align with several WHO criteria for NTDs: they are prevalent in tropical regions, disproportionately affect populations living in poverty, carry significant health and social consequences, and receive limited research funding and policy attention.⁵ Low socio-economic status was the single risk factor most consistently associated with SFIs in this review, supporting a parallel with the poverty-driven nature of NTDs. The geographic distribution is consistent with tropical endemicity, and the associated morbidity (psychological distress, alopecia) without reported mortality mirrors the predominantly morbidity-driven profile of NTDs. The paucity of antifungal susceptibility data and the apparent absence of national treatment guidelines further reflect the neglect characteristic of NTDs.

Importantly, the prevalence of SFIs far exceeds the prevalence of scabies (designated an NTD in 2017) by 20-fold or more in the reviewed studies, suggesting SFIs may warrant comparable recognition. However, the case for

A Systematic Review of Superficial Fungal Infections in Nigerian Children and a Case for Recognition as Neglected Tropical Diseases

NTD designation is weakened by the absence of school absenteeism data, disability-adjusted life year (DALY) estimates, and cost-burden analyses, which are essential for objective comparison with already-designated NTDs.⁶ Accordingly, while formal WHO NTD classification is not currently supported by the available evidence, SFIs in Nigerian children may reasonably be described as neglected dermatologic infections of poverty.

Methodological limitations and research gaps

All reviewed studies were cross-sectional studies, and no longitudinal data were presented. So, recurrence rates, chronicity, and long-term outcomes like psychosocial impact could not be studied. The predominance of school-based sampling limited random sampling methods, and the absence of a standardised diagnostic framework introduced risks of selection bias and comparability. Heavy reliance on clinical diagnosis, with only limited laboratory confirmation, further undermines data quality.

There was also an absence of nationally pooled estimates from a nationwide prevalence study across the six geopolitical zones, no cost-of-illness or DALYs studies, and no antifungal resistance surveillance. These gaps limit precise burden estimation. The overall quality of the studies was moderate to low due to poor randomisation, inconsistencies in the results with wide variations among the studies, and indirectness in assessing risk factors.

Future research recommendations

To remedy some of the research gaps highlighted, it will be prudent to conduct multi-centre nationwide prevalence surveys to establish national representative data on SFIs in Nigerian children. There should be standardised diagnostic criteria (clinical + microscopy + culture) to have uniform criteria for recruitment of study subjects. Establishing a fungal pathogen surveillance registry will improve the understanding of SFIs' epidemiology in Nigeria, as well as facilitate the investigation of antifungal resistance trends. Longitudinal

cohort studies should be performed to enable studying the determinants and outcomes of SFIs in Nigerian children. Studies should be conducted to evaluate school-based screening programs targeting SFIs, given the preponderance of these infections in the school-aged group. Cost-benefit and cost-effectiveness analyses of mass treatment or screening can be conducted to underscore the economic importance of SFIs as potential NTDs.

Policy recommendations

Nigerian educational and health policies should integrate SFIs into the school health programs to improve the control of these conditions. There should be health policies to include SFI surveillance within Nigeria's NTD strategic framework. This will help to leverage existing systems to provide a holistic approach to the treatment and control of childhood SFIs. In addition, policies should be developed on national treatment guidelines specific to paediatric populations, for more effective therapies. General health promotion policies should include hygiene education campaigns targeting primary schools. This will significantly impact the reduction of the prevalence of SFIs, which have been associated with poor hygiene practices. Lastly, national drug procurement policies should be directed at providing essential antifungal medications for treating SFIs in order to improve treatment access and reduce the number of children with untreated infections.

Conclusion

Superficial fungal infections, as typified by *tinea capitis*, are quite common among Nigerian children. Although they are not classically listed under WHO's list of NTDs, these SFIs share several features with known neglected diseases due to the high disease burden in low socio-economic classes, association with poor sanitation and overcrowding and limited surveillance and health policy attention. Despite this wide prevalence, SFIs among Nigerian children receive limited national public health

prioritisation, and there is inadequate aggregated data at the national level. Therefore, a call to designate SFIs in children as neglected dermatologic tropical diseases may draw attention and increase awareness of this underserved population.

Authors' Contributions: MHU conceived and designed the study and did the literature review, data analysis and drafting of the manuscript. MTT revised the draft manuscript for sound intellectual content. Both authors approved the final version of the manuscript.

Conflicts of Interest: None declared.

Financial support: The authors did not receive financial support for the research or for the preparation of the manuscript.

Accepted for publication: 3rd September 2026.

References

- Kelly BP. Superficial fungal infections. *Pediatr Rev* 2012;33:e22.
- World Health Organisation. Epidemiology and management of common diseases in children in developing countries. Geneva: World Health Organisation; 2005.
- Seebacher C, Bouchara JP, Mignon B. Updates on the epidemiology of dermatophyte infections. *Mycopathologia* 2008;166: 335–352.
- David OM, Ogunleye O, Adeola PO, Olawusi AC, Owabumoye JB, Ogidi C, *et al*. Dermatophytes and dermatophytoses acquisition among school-age children in Cameroon, Ivory Coast, Mali and Nigeria: A systematic review. *Caribb J Sci Technol* 2022;10(2):17–23.
- Jayashankar CA, Kumar BVP, Kandi V, Girish N, Sanjana K, Dharshini D, *et al*. Neglected tropical diseases: A comprehensive review. *Cureus* 2024;16(2):e53933.
- World Health Organisation. Summary of the Global Report on Neglected Tropical Diseases 2025. *Wkly Epidemiol Rec* 2026;101(3):5–12.
- Anosike J, Keke I, Uwaezuoke J, Anozie J, Obiukwu C, Nwoke B, *et al*. Prevalence and distribution of ringworm infections in primary school children in parts of Eastern Nigeria. *J Appl Sci Environ Manag* 2005;9(3):21–5.
<https://doi.org/10.4314/jasem.v9i3.17347>
- Enendu NE, Ibe SN. Prevalence of Tinea capitis among primary school pupils in Uli, Anambra State, Nigeria. *Afr J Appl Zool Environ Biol* [Internet]. 2005;7:1–4.
- Nweze EI, Okafor JI. Prevalence of Dermatophytic Fungal Infections in Children: A Recent Study in Anambra State, Nigeria. *Mycopathologia* 2005;160(3):239–43.
<https://doi.org/10.1007/s11046-005-0124-0>.
- Ogunbiyi AO, Owoaje E, Ndahi A. Prevalence of Skin Disorders in School Children in Ibadan, Nigeria. *Paediatr Dermatol* 2005;22(1):6–10.
- Uneke CJ, Ngwu BA, Egemba OS. Tinea Capitis And Pityriasis Versicolor Infections Among School Children in South-Eastern Nigeria: The Public Health Implications. *Internet J Dermatol* 2006;4(2). <https://doi.org/10.5580/2503>.
- Mbata T, Nwajagu C. Dermatophytes and Other Fungi Associated with Scalp Hair of Nursery and Primary School Children in Awka, Nigeria. *Internet J Microbiol* 2006;3(2).
- Oyedemi OA, Onayemi O, Oyedemi GA, Oyelami O, Aladekomo TA, Owa JA. Prevalence and pattern of skin infections and infestations among primary school pupils in Ijeshaland. *Niger J Paediatr* 2006;33(1):13–7.
- Ngwogu AC, Otokunfor TV. Epidemiology of dermatophytoses in a rural community in Eastern Nigeria and review of literature from Africa. *Mycopathologia* 2007;164(4):149–58.
<https://doi.org/10.1007/s11046-007-9038-3>
- Ayanbimpe GM, Taghir H, Diya A, Wapwera S. Tinea capitis among primary school children in some parts of central Nigeria. *Mycoses* 2008;51(4):336–40.
<https://doi.org/10.1111/j.1439-0507.2007.01476.x>
- Emele FE, Oyeka CA. Tinea capitis among primary school children in Anambra State of Nigeria. *Mycoses* 2008;51(6):536–41.
<https://doi.org/10.1111/j.1439-0507.2008.01507.x>.

17. Nweze EI. Dermatophytosis among children of Fulani/Hausa herdsman living in southeastern Nigeria. *Rev Iberoam Micol* 2010;27(4):191–4.
<https://doi.org/10.1016/j.riam.2010.06.003>
18. Sanuth HA, Efuntoye MO. Distribution and microbiological characterisation of dermatophyte infection among primary school children in Ago-Iwoye, Ogun State, Nigeria. *Researcher* 2010;2(6):95–9.
19. Amoran OE, Runsewe-Abiodun OO, Mautin AO, Amoran IO. Determinants of dermatological disorders among school children in Sagamu, Nigeria. *Educ Res* 2011;2(12):1743–8.
20. Okafor O, Akinbami F, Orimadegun A, Okafor C, Ogunbiyi A. Prevalence of dermatological lesions in hospitalised children at the University College Hospital, Ibadan, Nigeria. *Niger J Clin Pract* 2011;14(3):287.
<https://doi.org/10.4103/1119-3077.86769>.
21. Enemuor SC, Amedu AS. Prevalence of superficial mycoses in primary school children in Anyigba, Kogi State, Nigeria. *Afr J Microbiol Res* 2009;3(2):62–5.
22. Ayanlowo O, Akinkugbe A, Oladele R, Balogun M. Prevalence of Tinea capitis infection among primary school children in a rural setting in south-west Nigeria. *J Public Health Afr* 2014;5(1).
<https://doi.org/10.4081/jphia.2014.349>.
23. Oke OO, Onayemi O, Olasode OA, Omisore AG, Oninla OA. The Prevalence and Pattern of Superficial Fungal Infections among School Children in Ile-Ife, South-Western Nigeria. *Dermatol Res Pract* 2014;2014:1–7.
<https://doi.org/10.1155/2014/842917>.
24. Ogbu CC, Okwelogu IS, Umeh AC. Prevalence of superficial fungal infections among primary school pupils in Awka South. *J Mycol Res* 2015;2(1):15–22.
25. Dogo J, Afegbua SL, Dung EC. Prevalence of Tinea Capitis among School Children in Nok Community of Kaduna State, Nigeria. *J Pathog* 2016;2016:1–6.
<https://doi.org/10.1155/2016/9601717>.
26. Ahmed AA. Prevalence of Tinea Capitis among School Children in Kudan, Kaduna State. *Int J Sci Glob Sustain* [Internet]. 2017;4(1):98–107.
27. Afolabi OT, Oninla O, Fehintola F. Tinea capitis: A tropical disease of hygienic concern among primary school children in an urban community in Nigeria. *J Public Health Epidemiol* 2018;10(9):313–9.
<https://doi.org/10.5897/JPHE2018.1050>.
28. Damen DJG, Cosmas U, Iyala S. Distribution of Tinea capitis among primary school pupils in a local government area of Plateau State, Nigeria. *Eur J Biomed Pharm Sci* 2018;5(2):134–6.
29. Adesiji YO, Omolade BF, Aderibigbe IA, Ogungbe OV, Adefioye OA, Adedokun SA, et al. Prevalence of Tinea Capitis among Children in Osogbo, Nigeria, and the Associated Risk Factors. *Diseases* 2019;7(1):13.
<https://doi.org/10.3390/diseases7010013>.
30. Anya TC, Sunday BC. Prevalence of dermatophytic fungal infections among primary school children in Aba North, Abia State, Nigeria. *Int J Res Innov Appl Sci*. 2019;4(8):31–4.
31. Altraide DD, Alex-Hart BA. Prevalence and pattern of skin diseases among school-age children at the University of Port Harcourt Teaching Hospital, Nigeria: a hospital-based study. *Asian J Res Infect Dis* 2020;3(4):29–36.
<https://doi.org/10.9734/ajrid/2020/v3i430134>.
32. Odeigah L, Rotifa S, Shittu R, Mutalub Y. Prevalence and risk factors of superficial fungal infections among primary school pupils in Ilorin, North Central Nigeria. *Ann Afr Med Res* 2020;3(100):23–7.
33. Ayodele EH, Charles N, Abayomi F. Prevalence, identification and antifungal susceptibility of dermatophytes causing Tinea capitis in a locality of North Central Nigeria. *Afr J Infect Dis* 2021;15(1):1–9.
<https://doi.org/10.21010/ajid.v15i1.1>.
34. Ezomike NE, Ikefuna AN, Onyekonwu CL, Ubesie AC, Ojinmah UR, Ibe BC. Epidemiology and pattern of superficial fungal infections among primary school children in Enugu, south-east Nigeria. *Malawi Med J* 2021;33(1):21–7.
35. Olarinoye GM, Katibi OS, Apeh AJ, Fayemiwo SA, Ogunbiyi AO, George AO. Tinea capitis: aetiology, clinical features, and associated sociodemographic factors

- among school pupils in Ilorin, Kwara State, Nigeria. *Niger J Dermatol* 2021;11(3):36–47.
36. Onubogu CU, Ushie SN, Aghanya IN, Ugochukwu EF. Superficial fungal infections among school-age children: does prevalence and pattern differ between commercial and agrarian communities? *Asian J Res Infect Dis* 2022;11(2):22–32. <https://doi.org/10.9734/ajrid/2022/v11i2214>.
37. Ukhureigbe OM, Ajoseh SO, Adesemoye ET, Imanyikwa OI, Onawola OIE. Characterisation of dermatophytes isolated from primary school children in parts of some selected Southwest States of Nigeria. *Dutse J Pure Appl Sci* 2022;8(3a):186–96. Okpo EA, Andy IE, John GE, Chinyeaka RC. Epidemiology of dermatophytes among primary school children in Calabar, Nigeria. *Christ J Glob Health* 2024;11(1):84–90. <https://doi.org/10.15566/cjgh.v11i1.851>.
38. Abdu M, Musa FM, Yahya H. Prevalence of Tinea Capitis among Children Attending Some Islamiyya Schools in Kaduna Metropolis, Kaduna State, Nigeria. *Sahel J Life Sci FUDMA* 2025;3(4):483–90. <https://doi.org/10.33003/sajols-2025-0304-44>.